

PERSPECTIVE

TTV viral load as a predictor of antibody response to SARS COV-2 vaccination



Daniele Focosi, MD, PhD, MSc,^a Andreina Baj, MD, PhD,^b
Lorenzo Azzi, MD, PhD,^b Federica Novazzi, PhD,^b and
Fabrizio Maggi, MD, PhD^c

From the ^aNorth-Western Tuscany Blood Bank, Pisa University Hospital, Pisa, Italy; ^bDepartment of Medicine and Surgery, University of Insubria; and the ^cNational Institute for Infectious Diseases “L. Spallanzani”, Roma, Italy.

KEYWORDS:

torquetenovirus;
TTV;
COVID-19;
vaccine;
immune response

The measure of torquetenovirus (TTV) viremia is widely recognized as an optimal biomarker of an individual immune status. In the context of COVID-19, the predictive role of TTV load with regard to vaccine response has also been demonstrated, suggesting other intriguing applications for this widespread anellovirus.

J Heart Lung Transplant 2023;42:143–144

© 2022 International Society for Heart and Lung Transplantation. All rights reserved.

Torquetenovirus (TTV), a widespread anellovirus recognized as the main component of the healthy human virome, displays viremia that is highly susceptible to variations in immune competence. In general, higher viremias are associated with poor immune competence (e.g., causing opportunistic infections), while lower viremias are associated with hyperactive immune system (e.g., causing allograft rejection). In COVID-19, the applications of TTV monitoring is dual.

The first aspect is the correlation between the TTV loads and prognosis, which is rather controversial. Querido et al reported a kidney transplant recipient who had a TTV load of 5.6 log₁₀ copies/ml, which increased to 7.9 log₁₀ copies/ml on day 43 after the diagnosis of SARS-CoV-2 infection, when the patient seroreverted.¹ Froque et al reported that COVID-19 patients with detectable TTV DNA had an increased risk of subsequently developing infectious events (HR 9.28) and a trend ($p = 0.05$) toward higher TTV DNA area under a curve between days 7 and 17 after intensive care unit (ICU) admission in patients who died, as compared to survivors.² In line with the fact that the most severe

presentations of COVID-19 are due to an exaggerated immune response, Solis et al reported that the rate of severe cases was higher in patients with low TTV DNA load in plasma, considering a threshold of 700 copies/ml. In severe patients, SARS-CoV-2 viremia positivity rates were higher than those in mild-moderate cases at any time point. When combined, TTV DNA load and SARS-CoV-2 viremia allowed to predict the outcome of COVID-19 infection, with a higher risk (HR = 12.4) of ICU admission in patients with low TTV DNA load and positive SARS-CoV-2 viremia.³ Stincarelli et al reported that serum TTV DNA seems unrelated to COVID-19 severity, while TTV miRNA tth8 seems related.⁴ At the other end of the spectrum, Emmel et al found that the 100-day survival rate in cancer patients who died from COVID-19 was significantly lower in the TTV-positive group than in the TTV-negative group ($p = 0.0475$), and that in the cancer TTV-positive group, those who died also had a higher load of TTV than those who did not die ($p = 0.0097$).⁵

Saliva is an alternative (more easily accessible than blood) biological sample for PCR methods: Mendes-Correa et al reported that the saliva TTV and nasal-oro-pharyngeal SARS-CoV-2 loads were correlated ($p = 0.008$), and the TTV level decreased as symptoms resolved in the SARS-CoV-2 infected group ($p = 0.028$), but remained unchanged in the SARS-CoV-2 negative controls.⁶

Reprint requests: Daniele Focosi, MD, PhD, MSc, Department of Translational Research, Pisa University Hospital, via Paradisa 2, 56124 Pisa, Italy. Telephone/Fax: +39-050-996541.

E-mail address: daniele.focosi@gmail.com

A second, more relevant, aspect is the predictive role of TTV load with regard to vaccine responses. Healthy individuals with a median baseline TTV load of 3.8 log₁₀ copies/ml have an efficient immune response to the influenza vaccine 30 days after vaccination; more than 80% of healthy individuals with a median baseline TTV load of 4.1 log₁₀ copies/ml have a response to hepatitis B vaccine when measured at day 90 after vaccination.⁷ In lung transplantation, Hoek et al showed an odds ratio (OR) of 0.54 ($p < 0.0001$) for response to the first mRNA-1273 vaccine dose,⁸ while Gallais et al showed an adjusted OR of 17.8 ($p = 0.001$) for pre-vaccine TTV viral load ≥ 6.2 log₁₀ copies/ml and response to third BNT162b2 vaccine dose among 173 SARS-CoV-2-naïve recipients.⁹ In kidney transplantation, Reindl-Schwaighofer et al showed OR = 0.92 for homologous (ChAdOx1) or heterologous third dose in 2-dose mRNA vaccine nonresponders.¹⁰ No studies have been reported so far investigating TTV loads and COVID-19 vaccine responses in liver or hematopoietic stem cell transplantations. We have found that healthcare workers with a baseline TTV load equal to or below 3.4 log₁₀ copies/ml of serum showed higher anti-Spike antibody levels when measured on day 30 after the first BNT162b2 vaccine dose ($p < 0.001$; data not shown).

Compared to other solid organ transplant recipients, lung transplant recipients receive higher levels of immunosuppression and accordingly have reduced COVID-19 vaccine responses.¹¹ These findings suggest a continuum where the higher the immunosuppression, the better the correlation between the basal TTV loads and vaccine-elicited antibody responses. Reza Hosseini et al suggested that TTV load can be used to identify the optimal moment for vaccination in patients under maintenance immunosuppression,¹² and evidences published so far are strengthening this intuition. Despite at this stage further and larger observational clinical trials are required to consolidate this hypothesis, TTV-driven vaccine boosts represent an intriguing opportunity to increase the response rate of immunocompromised patients. In this regard, integration of a pilot trial within the Horizon2020-funded “TTV Guide Tx” project (<https://www.ttv-guide.eu/>) should be warranted, to assess utility in association with further correlates of vaccine response. So far only advanced age,⁸ mycophenolate use,^{9,13} triple immunosuppression,¹⁰ and more advanced stage of organ failure¹³ are reliable predictors of vaccine responses, while there are discordant results for time since transplantation.^{8,10}

Author contributions

D.F. searched the literature and wrote the first draft. **A.B., L.A.,** and **F.N.** provided raw data for the HCW cohort. **F.M.** revised the manuscript.

Disclosure statement

The authors declare no conflict of interest related to this manuscript.

References

1. Querido S, Calça R, Weigert A, et al. Kinetics of torquetenovirus DNA load in a recent kidney transplant recipient with mild SARS-CoV-2 infection and a failed antibody response. *Transpl Infect Dis* 2021;23:e13524.
2. Forqué L, Albert E, Giménez E, et al. Monitoring of Torque Teno virus DNAemia in critically ill COVID-19 patients: may it help to predict clinical outcomes? *J Clin Virol* 2022;148:105082.
3. Solis M, Gallais F, Garnier-Kepka S, et al. Combining predictive markers for severe COVID-19: Torquetenovirus DNA load and SARS-CoV-2 RNAemia. *J Clin Virol* 2022;148:105120.
4. Stincarelli MA, Baj A, Guidotti B, et al. Plasma Torquetenovirus (TTV) microRNAs and severity of COVID-19. *Virol J* 2022;19:79.
5. Emmel V, Gama B, de Paula A, Ferreira G, Binato R, Abdelhay E. Can Torque teno virus be a predictor of SARS-CoV-2 disease progression in cancer patients? *J Infect Chemother: Official J Japan Society of Chemotherapy* 2022;28:1623-7.
6. Mendes-Correa MC, Tozetto-Mendoza TR, Freire WS, et al. Torquetenovirus in saliva: a potential biomarker for SARS-CoV-2 infection? *PLoS One* 2021;16:e0256357.
7. Maggi F, Tempestini E, Lanini L, et al. Blood levels of TT virus following immune stimulation with influenza or hepatitis B vaccine. *J Med Virol* 2005;75:358-65.
8. Hoek RA, Verschuuren EA, de Vries RD, et al. High torque tenovirus (TTV) load before first vaccine dose is associated with poor serological response to COVID-19 vaccination in lung transplant recipients. *J Heart Lung Transplant: Official Publication of the International Society for Heart Transplant* 2022;41:765-72.
9. Gallais F, Renaud-Picard B, Solis M, et al. Torque teno virus DNA load as a predictive marker of antibody response to a three-dose regimen of COVID-19 mRNA-based vaccine in lung transplant recipients. *J Heart Lung Transplant: Official Publication of the International Society for Heart Transplant* 2022.
10. Reindl-Schwaighofer R, Heinzel A, Mayrdorfer M, et al. Comparison of SARS-CoV-2 Antibody response 4 weeks after homologous vs heterologous third vaccine dose in kidney transplant recipients: a randomized clinical trial. *JAMA Intern. Med.* 2022;182:165-71.
11. Boyarsky BJ, Werbel WA, Avery RK, et al. Antibody response to 2-Dose SARS-CoV-2 mRNA vaccine series in solid organ transplant recipients. *JAMA* 2021;325:2204-6.
12. Reza Hosseini O, Nielsen SD. Torque-Teno virus to optimize timing of vaccination against severe acute respiratory syndrome coronavirus 2 in solid organ transplant recipients. *J Clin Virol* 2021;143:104967.
13. Tanner R, Starr N, Chan G, et al. Humoral response to SARS-CoV-2 adenovirus vector vaccination (ChAdOx1 nCoV-19 [AZD1222]) in heart transplant recipients aged 18 to 70 years of age. *J Heart Lung Transplant: Official Publication of the International Society for Heart Transplant* 2022;41:492-500.